

## **Patient Information** Last Name:\_\_ First Name: MI: Sex:□M □F \_\_\_SSN: \_\_ DOB (mm/dd/yyyy):\_\_\_ Parent/Guardian (if applicable):\_ (REQUIRED if patient is <18 years old at time of sample collection) \_\_ City:\_\_ Phone: State: Zip: ICD-10 Diagnosis Code(s) Primary ICD-10 Code (REQUIRED) Additional ICD-10 Code(s) (OPTIONAL) **Billing Information BILL:** □Insurance\* □Laboratory □Patient/Self-Pay □Provider Account \*Please provide details below AND attach a front/back copy of insurance card(s). Insurance Carrier: Policy #: **Provider/Account Information** Account Name/Address: Phone: Fax: **Ordering Provider Selection & Signature (REQUIRED)** Clearly select the ordering provider from list below (circle or ✓). If provider is not listed, please provide their FULL NAME AND NPI: I certify the ordered test(s) is/are reasonable and medically necessary for the diagnosis, care, and/or treatment of this patient's condition as documented in the medical record. If this patient is <18 years old at the time of testing, the parent/guardian has been informed that they will be responsible for any costs associated with testing. †Informed consent for genetic testing: I have read and understand the genetic consent requirement on page two and acknowledge I have obtained consent from my patient. Name:\_ **Sample Collection Information** Date (mm/dd/yyyy): \_\_\_ $\square$ AM $\square$ PM \_\_\_\_ Sender Sample ID: \_ Patient ID: **MEDICARE ONLY** - HOSPITAL STATUS WHEN SAMPLE WAS COLLECTED: ☐ Hospital Inpatient ☐ Hospital Outpatient ☐ Non-Hospital Patient Laboratory/Other Name/Address: Phone: Fax: ☐ Mail ☐ Fax ☐ No Results to Lab

IBD: Inflammatory bowel disease; UC: Ulcerative colitis; CD: Crohn's disease; IV: Intravenous; SC: Subcutaneous; TPMT: thiopurine methyltransferase; HCV: Hepatitis C; NASH: Non-alcoholic steatohepatitis

## All information requested is REQUIRED to process testing and/or submit insurance claims on your patient's behalf.

Prometheus Client Services | 888-423-5227 | 6:00 am - 4:30 pm Pacific Time

_	Select Test(s) To Be Performed									
	Therapeutic Drug Monitoring —									
ш	Anser® IFX (#315	50)								
	Simultaneously measure Select biologic and ro									
	☐ZYMFENTRA® SC (IFX-dyyb)	□AVSOLA® IV (IFX-axxq)	□REN	FLEXIS® IV -abda)						
	□INFLECTRA® IV (IFX-dyyb)	□REMICADE® I (infliximab)	V □Othe	er:						
	Anser® ADA (#3	170)								
	Simultaneously measures adalimumab (ADA) and antibody-to-ADA (ATA) levels.  Select biologic (optional):									
	☐HUMIRA® (adalimumab)	□Other: —								
	Anser® VDZ (#3' Simultaneously measure		(VDZ) and an	tibody-to-VDZ (ATV)	) levels.					
	Simultaneously measures vedolizumab (VDZ) and antibody-to-VDZ (ATV) levels.  Anser® UST (#3190)									
	Simultaneously measures ustekinumab (UST) and antibody-to-UST (ATU) levels.  Select biologic (optional):									
	STELARA* Other: (ustekinumab)									
	Anser® RZB (#3	140)								
	Simultaneously measures risankizumab (RZB) and antibody-to-RZB (ATR) levels.									
	Anser Test	ing - Biologic I	Dosing Data	(optional)						
	Date of Last Dose	Curre		Current						
	(mm/dd/yyyy)	Dos	□mg		days					
			□mg/kg		weeks					
	RiskImmune®† (#3600) Assesses genetic predisposition to immunogenicity to anti-TNF therapy.									
Ш	TPMT Genetics  Aids in individualized th		based on pati	ent TPMT genotype.						
	TPMT Enzyme (#3320) Phenotyping test that aids in individualized thiopurine dosing.									
	Thiopurine Metabolites (#3200) Aids in optimization of thiopurine dosing to achieve clinical response.									
		tabolites (#	‡3200)	-						
	Aids in optimization of	tabolites (#	#3200) g to achieve c	linical response.						
	Aids in optimization of	tabolites (#	#3200) g to achieve c	linical response.	mg/day					
	Aids in optimization of	tabolites (# thiopurine dosing nt Thiopurine	#3200) g to achieve c  Dosing (optimg/day	linical response.	mg/day					
	Aids in optimization of  Curre  5-MP mg/day	tabolites (# thiopurine dosing int Thiopurine   AZA  — Celiac &	#3200) g to achieve c  Dosing (optimg/day	linical response.	ng/day					
	Curre 6-MPmg/day  Celiac PLUS <sup>†</sup> (# Includes Celiac Geneti	tabolites (# thiopurine dosing int Thiopurine AZA — Celiac & 6360) cs and Celiac Se	#3200) g to achieve c  Dosing (opti  mg/day  Liver —	linical response.	ng/day					
	Curre  S-MP mg/day  Celiac PLUS <sup>†</sup> (#	tabolites (# thiopurine dosing nt Thiopurine	#3200) g to achieve c  Dosing (opti  mg/day  Liver —  erology.	linical response.  ional)  Otherr	ng/day					
	Curre  S-MPmg/day  Celiac PLUS <sup>†</sup> (# Includes Celiac Geneti	tabolites (# thiopurine dosing Int Thiopurine   Int Thiop	#3200) g to achieve c  Dosing (opti  mg/day  Liver —  erology.	inical response.  Otherr	ng/day					
	Celiac PLUS <sup>†</sup> (# Includes Celiac Genetic Genetic profile to ai ONLY Celiac Serole Antibody profile to — ONLY the followin	tabolites (# thiopurine dosing Int Thiopurine   Int Thiop	#3200) g to achieve composing (option of the composition of the compos	inical response.  Otherr  Constantr	ng/day					
	Celiac PLUS <sup>†</sup> (# Includes Celiac Genetic Genetic profile to ai ONLY Celiac Serole Antibody profile to	tabolites (# thiopurine dosing Int Thiopurine   Int Thiop	#3200) g to achieve component(s) of gA (tTG IgA)  #3200) g to achieve component(s) of gA (tTG IgA)  #3200) g to achieve component(s) of gA (tTG IgA)  #3200)	inical response.  ional)  Other r  cee.  of Celiac Serology — (#1405)	ng/day					
	Celiac PLUS <sup>†</sup> (# Includes Celiac Genetic PonLy Celiac Serold Antibody profile to  ONLY Celiac Serold Antibody profile to  Anti-Human Tissue Tr.  Anti-Endomysial IgA (#	tabolites (# thiopurine dosing Int Thiopurine   Int Thiop	#3200) g to achieve component(s) of gA (tTG IgA) Total set	inical response.  ional)  Otherr  cse. of Celiac Serology — (#1405) rum IgA (#1605)	ng/day					
	Curre  Curre  Curre  S-MP mg/day  Celiac PLUS <sup>†</sup> (# Includes Celiac Geneti ONLY Celiac Geneti ONLY Celiac Serole Antibody profile to ONLY the followin Anti-Human Tissue Tr. Anti-Endomysial IgA (# DGP IgA (#1255)  FIBRO Spect® I	tabolites (#thiopurine dosing int Thiopurine dosing designated colors and diagnosis of a dosignated colors dosignated colors dosignated colors dosignated colors dosignated colors dosignated dosignated colors dosignated dosignated colors dosignated dosig	#3200) g to achieve composing (optime mg/day)  Liver — erology.  eliac disease omponent(s) of gA (tTG IgA)  by Total set  or HCV patient	inical response.  ional)  Otherr  cse. of Celiac Serology — (#1405) rum IgA (#1605)	ng/day					
	Curre	tabolites (#thiopurine dosing int Thiopurine dosing designated colors and diagnosis of a dosignated colors dosignated colors dosignated colors dosignated colors dosignated colors dosignated dosignated colors dosignated dosignated colors dosignated dosig	#3200) g to achieve composing (optime mg/day)  Liver — erology. eliac disease component(s) of gA (tTG IgA) (so IT)  or HCV patient (so IT)  or HCV patient (so IT)  or NASH patient (so IT)	inical response.  ional)  Otherr  cse. of Celiac Serology — (#1405) rum IgA (#1605)	ng/day					
	Curre	tabolites (#thiopurine desired thiopurine desired t	#3200) g to achieve composing (optime mg/day)  Liver — erology.  eliac disease opponent(s) of gA (tTG IgA) of Total set of	linical response.  ional)  Otherr  se. of Celiac Serology — (#1405) rum IgA (#1605)  ints.  ents.	above.					

- Page 1 - DX.1002v12 (09/2025)

<sup>-</sup> Specimen collection requirements on page 2 -

<sup>&</sup>lt;sup>†</sup>Acknowledgment of informed consent for genetic testing required.

\*Failure to provide the required data, at the time of sample collection, may result in reporting delay/cancellation.

Specimen Collection and Handling Procedures								
Test Ordered	Turnaround Time <sup>a</sup>	Transport Kit and Storage Conditions	Specimen Type and Volume	Specimen Collection Tube	Specimen Stability			
Anser* ADA, IFX, RZB, UST, VDZ	3 days	Ambient or cold pack <u>Do not freeze</u>	2.0 mL Serum (0.5 mL for Peds)	Serum Separator Tube or Red-Top Tube	Room temp: 14 days Refrigerated: 14 days			
Celiac PLUS <sup>†</sup>	4 days	Ambient or cold pack <u>Do not freeze</u>	2.0 mL Serum <b>AND</b> 2.0 mL Whole Blood	Serum Separator Tube or Red-Top Tube AND EDTA/Lavender-Top Tube	Room temp: 7 days Refrigerated: 30 days			
Celiac Genetics†	4 days	Ambient or cold pack <u>Do not freeze</u>	2.0 mL Whole Blood	EDTA/Lavender-Top Tube	Room temp: 7 days Refrigerated: 30 days			
Celiac Serology (including all components)	4 days	Ambient or cold pack <b>Do not freeze</b>	2.0 mL Serum (0.5 mL for Peds)	Serum Separator Tube or Red-Top Tube	Room temp: 7 days Refrigerated: 30 days			
FIBRO <i>Spect</i> ® HCV, NASH	7 days	Ambient or cold pack <u>Do not freeze</u>	2.0 mL Serum (0.5 mL for Peds)	Serum Separator Tube or Red-Top Tube	Room temp: 7 days Refrigerated: 30 days			
IBD sgi Diagnostic*†	4 days	Ambient or cold pack <b>Do not freeze</b>	2.0 mL Serum <b>AND</b> 2.0 mL Whole Blood	Serum Separator Tube or Red-Top Tube AND EDTA/Lavender-Top Tube	Room temp: 7 days Refrigerated: 21 days			
RiskImmune*†	4 days	Ambient or cold pack. <u>Do not freeze</u>	2.0 mL Whole Blood	EDTA/Lavender-Top Tube	Room temp: 10 days Refrigerated: 30 days			
Thiopurine Metabolites	3 days	Cold pack preferred Do not freeze	5.0 mL Whole Blood	EDTA/Lavender-Top Tube	Room temp: 24 hours Refrigerated: 8 days			
TPMT Enzyme	3 days	Cold pack preferred Do not freeze	5.0 mL Whole Blood	EDTA/Lavender-Top Tube	Room temp: 24 hours Refrigerated: 8 days			
TPMT Genetics <sup>†</sup>	4 days	Ambient or cold pack <u>Do not freeze</u>	2.0 mL Whole Blood	EDTA/Lavender-Top Tube	Room temp: 10 days Refrigerated: 30 days			

<sup>&</sup>lt;sup>a</sup>Business days from date of receipt.

SHIPPING INSTRUCTIONS: Prometheus has an agreement with FedEx® for express overnight delivery within the United States and Canada. Please call FedEx at 1-800-GoFedEx (463-3339) to schedule a pickup. FedEx will pick up your specimens and ship them to Prometheus Laboratories Inc. in San Diego at no expense to you. Prometheus will provide specimen transportation kits upon request. NOTE: Multiple specimens may be shipped in a single transportation kit. For more information, call Client Services at 888-423-5227, or go to prometheuslabs.com.

## **ACKNOWLEDGMENT OF INFORMED CONSENT FOR GENETIC TESTING**

I warrant that this test was ordered and that I have obtained the appropriate prior written consent. This written consent was signed by the person who is the subject of the test (or if that person lacks capacity to consent, signed by the person authorized to consent for that person) and includes the following (unless certain that the following information is not required by the state in which I practice):

- 1. The purpose of this test is to determine if the patient may have a variant in the gene(s) being tested, which has been found to be associated with this condition.
- 2. This test will only test for this specific condition and will not detect ALL possible variants within this gene, nor will it detect variants in other genes.
- 3. Prior to my signature of the written consent, a qualified healthcare professional discussed with the patient the genetic test ordered and described the steps involved in the test, the constraints of the procedure, and its accuracy.
- 4. My patient was advised by a qualified healthcare professional of the risks and benefits of genetic testing, and advised of the significance of a positive and negative result.
- 5. The patient understands that a positive test result is an indication that the patient may be predisposed to, or have, the condition listed.
- 6. The patient understands that the test may fail, that the results may be non-informative or not predictive for his/her case, and that the test may reveal information that is unrelated to their intended purpose.
- 7. No unauthorized test is performed on specimens.
- 8. NY-state specimens will be destroyed within 60 days after collection when no longer required for clinical purposes.
- 9. The informed consent does not authorize the use or release of any other identified medical information unrelated to this genetic test
- 10. The patient understood that he/she could see professional genetic counseling prior to undergoing the testing procedure, and received written information identifying a genetic counselor or medical geneticist by his/her treating provider.

Prometheus tests are laboratory-developed tests that were developed, and analytically and clinically validated by Prometheus Laboratories Inc. under federal Clinical Laboratory Improvement Amendments (CLIA) guidelines, and are performed exclusively in our high complexity CLIA certified and College of American Pathologists accredited clinical laboratory. As laboratory developed tests, they have not been cleared or approved by the US FDA. The tests may be covered by one or more US pending or issued patents - see prometheuslabs.com/patents. Prometheus, Anser, FIBROSpect, IBD sgi Diagnostic and RiskImmune are registered trademarks of Prometheus Laboratories Inc, San Diego, California. All other trademarks or service marks are the property of their respective owners. ©2025 Prometheus Laboratories Inc. All rights reserved.

<sup>†</sup>Acknowledgment of informed consent for genetic testing required.

Specimens should be labeled with 2 identifiers and date of collection. Examples of acceptable identifiers include, but are not limited to: patient name, date of birth, hospital number, requisition, accession, or unique random number. Unlabeled specimens will not be accepted for testing.

SHIPPING INSTRUCTIONS: Prometheus has an agreement with FedEx\* for express overnight delivery within the United States and Canada. Please call FedEx at 1-800-GoFedEx (463-3339)